



***In vivo* genotoxicity of 2-alkylcyclobutanones in liver cells from rats fed with irradiated cocoa butter using flow cytometry.**

Martins¹ R., Vieira¹ D. P., Carvalho¹ L R., Barbezan¹ A. B., Villavicencio¹., A.L.C.H.

¹ Instituto de Pesquisas Energéticas e Nucleares (IPEN-CNEN - SP)
Av. Professor Lineu Prestes 2242
05508-000 São Paulo, SP, Brazil
regivip@gmail.com

ABSTRACT

Food irradiation proves to be an effective technique of eliminating some pathogens from food and this has gained significant attention to its potential for food safety. Since 1990, studies on the toxicological safety of 2-Alkylcyclobutanones have been conducted extensively. 2-Alkylcyclobutanones are unique radiolytic products generated by the radiation-induced breakage of triglycerides in food, 2-dodecylcyclobutanone (2-dDCB) and 2-tetradecylcyclobutanone (2-tDCB) are the predominant compounds detected. Despite studies showing non-genotoxicity of 2-ACBs (2-Alkylcyclobutanones), the results are conflicting and therefore we continue the studies in order to confirm the compounds are safety for human health. *In vivo* micronucleus test were performed to verify the 2-ACBs genotoxic effects in hepatic cells using flow cytometry. We used cocoa butter irradiated with 20 kGy. A group with animals was treated with daily intake of irradiated cocoa butter, synthesized 2-dDCB and 2-tDCB for one month. Hepatic cells were selected for genotoxicity analysis due to the liver importance in the compounds metabolism. The improvement of the analytical techniques is important for the research future since the irradiation process is already consolidated. The results confirmed the safety of the food irradiation process, as they did not indicate the genotoxic potential of the samples.

Keywords: 2- Alkylcyclobutanones, Micronucleous test, liver.

1. INTRODUCTION

1.1 Food irradiation

Food irradiation has been demonstrated as an effective tool and gained great attention due to its potential for food safety assurance, reducing microbial load, insect pest control in agricultural products, extension of the period of shelf, budding inhibition [1].

The critical factor in this process is how to apply precise radiation doses to meet the requirement of the process without causing significant losses of sensory and nutritional values to foods [2].

However, there are some limitations as well: the formation of radiolytic compounds such as 2-alkylcyclobutanones that are derived from the irradiation of foods containing fat [3].

1.2 2-Alkylcyclobutanones formation


Since 1990, studies on the toxicological safety of 2-Alkylcyclobutanones have been conducted extensively [4]. 2-Alkylcyclobutanones are unique radiolytic products generated by the radiation-induced breakage of triglycerides in food, are exclusively found in irradiated lipid containing foods [5], 2-dodecylcyclobutanone from radiolysis of palmitic acid (2-dDCB) and 2-tetradecylcyclobutanone from stearic acid (2-tDCB), are the predominant compounds detected [6]. They have the same number of carbons (n) as their fatty acid precursors, with an alkyl chain of carbons in ring position 2 [7] Fig.1. and are thus considered unique markers of food irradiation [8].

The toxic potential of 2-ACBs has been previously investigated by a French–German research group in an EU interregional programme [9], in which the tumour-promoting potential of 2-tDCB and 2-(tetradec-5-enyl)-cyclobutanone (2-tDeCB) was reported [10]. Despite studies showing non-genotoxicity of 2-ACBs, the results are conflicting and therefore we continue the studies in order to confirm the compounds safety for human health. In vivo micronucleus test were performed to verify

the 2-ACBs genotoxic effects in hepatic cells using flow cytometry. However, further studies are required to resolve remaining uncertainties regarding the genotoxicity of 2-ACBs, because very few studies have been conducted to investigate the genotoxicity of pure 2-ACBs according to specific test guidelines [11].

Figure 1: Structure of the basic side chains of the four most common Alkylcyclobutanones found in foods containing lipids irradiated.

Irradiation Dose Indicators



2. MATERIAL AND METHODS

2.1 Samples

Cocoa butter samples containing 29% palmitic acid (2-Dodecylcyclobutanone precursor) and 32% stearic acid (2-Tetradecylcyclobutanone precursor) were used [13].

It was gamma-irradiated (20 kGy, dose rate 0.473 kGy/h) using a ^{60}Co source (IPEN Gammacell 220) at room temperature.

They were fluidized in a water bath at 38°C and administered orally daily to all animals 1 mL/kg/bw.

2.2 Chemicals

Type I collagenase (cas n.9001-12-1 sigma aldrich[®]) was solubilized (300 µg/ml) in hank's balanced salt solution (HBSS). Cyclophosphamide monohydrate (cp) (cas n.6055-19-2 sigma aldrich[®]) was solubilized in saline solution (NaCl 0,9%).

The synthesized compounds 2-alkylcyclobutanones (2-dodecylcyclobutanone, cas n.35493-46-0 e 2-tetradecylcyclobutanone, cas n.35493-47-1, both sigma aldrich[®]), were solubilized in vegetable soybean oil in a concentration of 4 mg/ml.

Cell suspensions were properly incubated with sytox[®] green (molecular probes, s7020) and ethidium monoazide bromide (ema, molecular probes, e1374) to label nuclear and micronuclear dna (sytox) and to discriminate nuclei from dead cells (ema).

2.3 Animals housing conditions

Groups of 12 male *wistar* rats were bred and maintained for two months at ipen animal facility under all principles of animal welfare (12/12 h day night cycles, room temperature: 22-23 °c, 44-65% relative atmospheric humidity, no more than three animals/cage), as specified by ipen ethical animal experimentation committee (process number ceua-ipen No.148/14), including food and water *ad libitum* availability.

Two animals were supplemented with daily oral dose of 1g of irradiated cocoa butter containing 2-alkylcyclobutanones, two animals were treated with daily oral dose of 4 mg / kg of 2-dodecylcyclobutanone and 2 animals with the same dose of 2-tetradecylcyclobutanone dissolved in oil for 5 days.

Positive controls for genotoxicity were represented by 2 animals injected intraperitoneally with cyclophosphamide (50 mg/kg/bw), 24 hours prior the experiment.

Negative controls were represented by 2 animals injected with 1 ml of vehicle control (NaCl 0,9 %) and more two animal without treatment and interferences, according to OECD standards, which are a unique tool for assessing the potential effects of chemicals on human health and the environment. All samples were performed in quadruplicates.

2.4 Flow cytometry

2.4.1. Hepatic cell isolation

Animals were euthanatized by exposure to CO₂ approximately 24 h following the last treatment. Livers were exposed and fragments of 3 to 4 mm³, were dissociated passing through syringe needle in HBSS with collagenase (200 µg/ml). Tissue pieces were incubated at 37 °C for 1 hour, with vigorous shaking every 15 minutes.

Suspensions were centrifuged (1500 rpm, 5 min, RT) and cell pellets were suspended in ammonium chloride (15.2 mM in water) to lyse erythrocytes and kept on ice for 5 minutes. Cells were centrifuged as described and suspended in ice-cold phosphate-buffered saline solution (PBS).

2.4.2. Flow cytometric analysis

Isolated liver cells from rats were plated (100 µL/well) in 96-well plates in quadruplicates. Briefly, cells were incubated with EMA (5 µg/mL) and exposed to blue led light (30 W) in an ice bath to photoactivation of dye.


Plates were centrifuged (1500 rpm, 15 min, RT) and cells were lysed with buffer [14].

3. RESULTS AND DISCUSSION

Body weight and food intake in each group were similar throughout the experimental period. We administered 2-dDCB or 2-tDCB to the rats orally for 5 days. Statistically significant increases were observed in the frequency of micronucleous (MNs) after doses of 4 mg/kg/bw day of 2-dDCB and 2-tDCB.


There were significant differences between the 2-dDCB or 2-tDCB groups and the vehicle control group. On the other hand, the group that ingested the irradiated cocoa butter, representing the amount naturally ingested by food, did not present significant amounts that induce genotoxicity (Graf.1). The chemical positive control (CP) clearly induced MNs (Fig.2).

Graphic 1: Percentage of MN (% of total events) of liver cells of rats.



CC (cellular control), NaCl (Negative control), Ciclophosphamide (Positive control), 2-Dodecylcyclobutanone, 2-Tetradecylcyclobutanone and 20 kGy irradiated cocoa butter. Bars: SEM (Standard error of means). (**): $p < 0,01$ (difference from negative control).

Figure 2: Representative Flow cytometry of experimental examples from group results.



(A) Cells control, (B) NaCl 0,9%; (C) Ciclophosphamide, (D) 2-Dodecylcyclobutanone, (E) 2-Tetradecylcyclobutanone and (F) 20 kGy Irradiated cocoa butter.

According to the literature, few *in vivo* studies have been performed to evaluate the genotoxicity of 2-ACBs.

In 2002, Horvatovich et.al [15], reported that less than 1% of 2-ACBs are excreted in feces and a small amount was detected in adipose tissue.

Raul et.al 2002[10], affirmed the promotion of colon cancer, causing the scientific community to perform more studies on the subject. Until then studies using micronuclei had not been performed.

Yamakage et.al 2014[11] performed the bone marrow micronucleus test and suggested the non-genotoxicity of 2-ACBs.

The last micronucleus test performed by Beom S.Song et.al 2018[16], uses doses of 250, 500, 1000 and 2000 mg/kg/bw and has results in line with Yamakage et.al 2014. But compared to the present article there is divergence in the results when it comes to micronucleus analysis in specific target organ cells.

Previous studies used bone marrow, while this study used cells extracted from liver tissue, which reported an increase in the number of MN.

The rat liver micronucleus test yields positive results with complex metabolic transforming substances, which negative results are obtained in bone marrow MN, metabolites produced in the liver do not reach the bone marrow. This technique using rat liver removes the problem of specificity by studying different target organs, according Cliet et al 1989 [17].

4. CONCLUSION

We conclude that the consume of 2-ACBs through food in this study represented by irradiated cocoa butter has no genotoxic potential.

Specific cell analyzes obtained satisfactory results indicating that they are more appropriate than bone marrow cell analyzes for certain compounds.

The in vivo micronucleus test by flow cytometry proved to be a satisfactory method for analyzing the genotoxicity of 2-Alkylcyclobutanones.

ACKNOWLEDGMENTS

Thanks to IPEN Gammacell irradiator for processing samples, CNEN, CNPq and FAPESP.

REFERENCES

- [1] SATIM, M. Food Irradiation a Guidebook, **Technomic Publishing Co.Inc**, Lancaster, PA, 1993.
- [2] FARKAS, J.; EHLERMANN, D.A.E. “Food irradiation”, **In:Encyclopedia of Food Safety. Foods, Materials, Techn Mohacsi-Farkasologies and Risks**, v.3, p. 178-186, 2014.
- [3] VARIYAR, P.S.; CHATTERJEE, S.; SAJILATA, M.G.; SINGHAL, R.S.; SHARMA,A. Natural existence of 2-alkylcyclobutanones. *Journal of Agricultural and Food Chemistry*, v. 56, p. 11817– 11823, 2008.
- [4] DELINCÉE, H.; POOL-ZABEL, B.L. Genotoxic properties of 2-dodecylcyclobutanone, a compound formed on irradiation of food containing fat. *Radiation Physics and Chemistry*, V.52, p.39–42, 1998.
- [5] LETELLIE, P.R.; NAWAR, W.W. 2-Alkylcyclobutanones from radiolysis of triglycerides. *Lipids*, v.7, p. 75–76, 1972.
- [6] CHAN, W.C.; YE, Y. LEUNGE,E.M.K. Rapid identification of γ -irradiated food by direct solvent extraction and liquid chromatography-tandem mass spectrometric analysis of 2-Dodecylcyclobutanone: Application in surveillance of irradiated food. **Food chemistry**, v.161, p. 312-316, 2014.
- [7] STEVENSON, M.H. Validation of the cyclobutanone protocol for detection of irradiated lipid containing foods by interlaboratory trial. In: MC MURRAY , C.H.; STEWART, E.M.; GRAY,R;

- PEARCE, J., (Eds.), *Detection Methods for Irradiated Foods – Current Status*. **Royal Society of Chemistry, Cambridge, UK**, p. 269–284, 1996.
- [8] NDIAYE B.; JAMET, G.; MIESCH, M.; HASSELMANN, C.; MARCHIONI, E. 2-Alkylcyclobutanones as markers for irradiated foodstuffs II. **The CEN (European Committee for Standardization)** method: field of application and limit of utilization, *Radiation Physics and Chemistry*, v. 55, p. 437–445, 1999.
- [9] BURNOUF, D.; DELINCÉE, H.; HARTWING, A.; MARCHIONI, E.; MIESCH, M.; RAUL F.; WERNER, D. Etude toxicologique transfrontalière destinée à évaluer le risque encouru lors de la consommation d'aliments gras irradiés – Toxikologische Untersuchung zur Risikobewertung beim Verzehr von bestrahlten fetthaltigen Lebensmitteln – Eine französisch-deutsche Studie im Grenzraum Oberrhein, *Rapport final d'étude Interreg II*, project No 3171, A summary of the report is available on Internet at: <http://www.iaea.org/programmes/rifa/icgfi/documents/summary-pres.s.pdf>, 2002.
- [10] RAUL, F.; GOSSÉ, F.; DELINCÉE, H.; HARTWING, A.; MARCHIONI, E.; MIESCH, M.; WERNER, D.; BURNOUF, D. Food-borne radiolytic compounds (2-alkylcyclobutanones) may promote experimental colon carcinogenesis. *Nutrition and Cancer*, v.44, p. 88–191, 2002.
- [11] YAMAKAGE, K.; SUL, H.; OHTA, R.; TYOIZUMI, T.; KAWAKAMI, K.; MATSUMOTO, H.; TAKAHASHI, T.; SASAKI, K.; IKEZUME, M.; NEGISHI, S.N.; IZUME, K.; TODORIKI, S.; TAKAHASHI, K.; FURUTA, M. Genotoxic potential and in vitro tumour-promoting potential of 2-dodecylcyclobutanone and 2-tetradecylcyclobutanone, two radiolytic products of fatty acids. *Mutation Research*, V.770, p. 94-104, 2014.
- [12] GADGIL, P.; HACHMEISTER, K.; SMITH, J.S.; KROPF, D.H. 2-Alkylcyclobutanones as Irradiation Dose Indicators in Irradiated Ground Beef Patties. *Journal of Agricultural and Food Chemistry*, v.50, p. 5746-5750, 2002.
- [13] WANG, H.; MALEKY, F. Effects of cocoa butter triacylglycerides and minor compounds on oil migration. *Food Research International*, v.106, p. 213-224, 2018.
- [14] BRYCE, S.M.; BEMIS, J.C.; AVLASEVICH, S. L.; DERTINGER, S.D. *In vitro* micronucleus assay scored by flow cytometry provides a comprehensive evaluation of cytogenetic damage and cytotoxicity, *Mutation Research*, v.630, p.78-91, 2007.

- [15] HORVATOVICH, P.; RAUL, F.; MIESCH, M.; BURNOUF, D.; DELINCÉE, H.; HARTWING, A.; WERNER, D.; MARCHIONI, E. Detection of 2-alkylcyclobutanones, markers for irradiated foods, in adipose tissues of animals fed with these substances. **Journal of Food Protection**, v.65, p. 1610–1613, 2002.
- [16] SONG, B.S.; KIM, Y.; JIN, Y.B.; KANG, I.J.; KIM, K.S.; PARK, J.K.; KIM, J.K.; PARK, H.Y.; JEONG, S.H. Toxicological evaluation of 2-Dodecylcyclobutanone, a unique radiolytic compound of palmitic acid, **Food and Chemical Toxicology**, v.121, p. 639-647, 2018.
- [17] CLIET, I.; FOURNIER, E.; MELCION, C.; CORDIER, A. In vivo micronucleus test using mouse hepatocytes, **Mutation Research**, v.216, p. 321-326, 1989.